

Connected Sums of Special Lagrangian Submanifolds

Dan A. Lee
Stanford University
dalee@math.stanford.edu

February 1, 2008

Abstract

Let M_1 and M_2 be special Lagrangian submanifolds of a compact Calabi-Yau manifold X that intersect transversely at a single point. We can then think of $M_1 \cup M_2$ as a singular special Lagrangian submanifold of X with a single isolated singularity. We investigate when we can regularize $M_1 \cup M_2$ in the following sense: There exists a family of Calabi-Yau structures X_α on X and a family of special Lagrangian submanifolds M_α of X_α such that M_α converges to $M_1 \cup M_2$ and X_α converges to the original Calabi-Yau structure on X . We prove that a regularization exists in two key cases: (1) when $\dim_{\mathbb{C}} X = 3$, $\text{Hol}(X) = \text{SU}(3)$, and $[M_1]$ is not a multiple of $[M_2]$ in $H_3(X)$, and (2) when X is a torus with $\dim_{\mathbb{C}} X \geq 3$, M_1 is flat, and the intersection of M_1 and M_2 satisfies a certain angle criterion. One can easily construct examples of the second case, and thus as a corollary we construct new examples of non-flat special Lagrangian submanifolds of Calabi-Yau tori.

1 Introduction

One of the fundamental problems in special Lagrangian geometry is to understand moduli spaces of special Lagrangian submanifolds (SLags). Much interest in this problem arises from the study of mirror symmetry since it is related to the SYZ Conjecture [19]. McLean's deformation theorem [17] together with some work by Hitchin [8] provide some understanding of these moduli spaces locally near nonsingular SLags, but in order to understand these moduli spaces globally, we need to understand singular SLags. Recently, some research has focused on the more modest goal of understanding SLags with isolated conical singularities. For example, see [10, 11, 12, 13]. In order to study SLags with isolated conical singularities, we need to know something about the SLag cones in $\mathbb{C}^n$ on which these singularities are modelled. Some researchers have undertaken the study of these SLag cones. See [7].

In this paper we restrict our attention to isolated conical singularities modelled on a very simple type of SLAG cone, namely, the union of two transversely intersecting SLAG planes in $\mathbb{C}^n$. By the work of Lawlor [15], we know that such a SLAG cone can be deformed through a family of nonsingular SLAGs in $\mathbb{C}^n$ if the two planes meet a certain angle criterion to be described later. (This criterion is always satisfied when $n \leq 3$.) This local regularization holds out hope that if we have a singular SLAG with this simple type of singularity, then it can be globally regularized, that is, it can be deformed through a family of nonsingular SLAGs. This simple type of singularity arises when a connected, immersed SLAG intersects itself, and when two embedded SLAGs intersect. The first case has already been treated by Yng-Ing Lee [16] who answered the question in the affirmative: A compact, connected, immersed SLAG with an isolated point of transverse self-intersection satisfying the angle criterion can be regularized. The second case is more difficult, and that is the case which we consider in this paper. Simply put, our problem is to try to regularize the union of two compact embedded SLAGs with an isolated point of transverse intersection satisfying the angle criterion. A problem related to ours has been solved by Butscher [1]: The union of two embedded SLAGs with boundary in $\mathbb{C}^n$ with $n \geq 3$ with an isolated point of transverse intersection satisfying the angle criterion can be regularized. In Butscher's paper, the regularization takes advantage of the freedom to deform the boundary of the singular SLAG. In our problem, we have no boundaries, and therefore we cannot use the added degrees of freedom. Indeed, our problem as stated above probably cannot be solved. We must introduce another degree of freedom, and we do this by deforming the Calabi-Yau structure of the ambient manifold.

Before we state our results, we recall some basic definitions and facts.

Definition. A **Calabi-Yau structure** (or CY structure) on a compact $2n$ -fold X is a 3-tuple (J, ω, Ω) such that J is a complex structure on X , ω is a Kähler form with respect to J , and Ω is a holomorphic $(n, 0)$ form with respect to J such that

$$\frac{\omega^n}{n!} = (-1)^{\frac{1}{2}n(n-1)} \left(\frac{i}{2}\right)^n \Omega \wedge \overline{\Omega} \quad (1)$$

It is a fact that $\text{Re } \Omega$ is a calibration with respect to the Kähler metric. We say that a submanifold M of X is **special Lagrangian** iff M is calibrated by $\text{Re } \Omega$.

The special Lagrangian condition on M is equivalent to the vanishing of both ω and $\text{Im } \Omega$ on M . If (X, J) admits a Calabi-Yau structure at all, then in each Kähler class there is a unique Kähler form ω such that (X, J, ω) admits a Calabi-Yau structure. (In this case, the Kähler metric corresponding to ω is Ricci-flat.) Also, if (X, J) admits a Calabi-Yau structure, then Ω is uniquely determined up to a complex constant. If we also choose ω , then the normalization (1) uniquely determines Ω up to a phase. Because of these facts, a choice of Calabi-Yau structure amounts to a choice of complex structure, a Kähler class, and a phase. For more general background on Calabi-Yau manifolds and special Lagrangian geometry, see [18, 9, 14, 6].

We now describe the angle criterion.

Definition. Given any two n -dimensional oriented linear subspaces η and ξ of $\mathbb{R}^{2n}$, there exist **characterizing angles** $\theta_1, \dots, \theta_n$, together with an orthonormal basis $e_1, \dots, e_{2n}$ of $\mathbb{R}^{2n}$, such that

$$0 \leq \theta_1 \leq \dots \leq \theta_{n-1} \leq \frac{\pi}{2} \text{ and } \theta_{n-1} \leq \theta_n \leq \pi - \theta_{n-1}$$

while

$$\eta = e_1 \wedge \dots \wedge e_n$$

and

$$\xi = [(\cos \theta_1)e_1 + (\sin \theta_1)e_{n+1}] \wedge \dots \wedge [(\cos \theta_n)e_n + (\sin \theta_n)e_{2n}].$$

We say that η and ξ satisfy the **angle criterion** iff the characterizing angles between η and $-\xi$ satisfy $\sum_{i=1}^n \theta_i = \pi$.

The Lawlor-Nance Angle Theorem states that a pair of oriented planes (η, ξ) is minimizing iff the characterizing angles between η and $-\xi$ satisfy $\sum_{i=1}^n \theta_i \geq \pi$. Therefore the angle criterion may be thought of as describing the “borderline case” of minimizing pairs of planes. See [6] for more on characterizing angles and the Angle Theorem.

We are now ready to state our main theorem.

Theorem 1 (Main Theorem). *Let M_1 and M_2 be two embedded special Lagrangian submanifolds of a Calabi-Yau manifold (X, J, ω, Ω) such that $n = \dim_{\mathbb{C}} X \geq 3$ and the holonomy of the Kähler metric is exactly $SU(n)$. Assume that M_1 and M_2 intersect transversely at a single point p such that the tangent cone of $M_1 \cup M_2$ at p satisfies the angle criterion. Further assume that $\text{Re}[H^{n-1,1}(X) \oplus H^{1,n-1}(X)]$ is not contained in the kernel of $\frac{[M_1]}{\text{Vol}(M_1)} - \frac{[M_2]}{\text{Vol}(M_2)}$, thought of as a functional on $H^n(X)$. Then there exists a family of Calabi-Yau structures $(J_\alpha, \omega_\alpha, \Omega_\alpha)$ on X converging to (J, ω, Ω) and a family of embedded submanifolds $M_\alpha \subset X$ converging to $M_1 \cup M_2$ such that M_α is special Lagrangian in $(X, J_\alpha, \omega_\alpha, \Omega_\alpha)$.*

When $n = 3$, the theorem reduces to the following nice result.

Corollary 2. *Let M_1 and M_2 be two embedded special Lagrangian submanifolds of a Calabi-Yau manifold (X, J, ω, Ω) such that $\dim_{\mathbb{C}} X = 3$ and the holonomy of the Kähler metric is exactly $SU(3)$. Assume that M_1 and M_2 intersect transversely at a single point and that $[M_1]$ is not a multiple of $[M_2]$ in $H_n(X)$. Then there exists a family of Calabi-Yau structures $(J_\alpha, \omega_\alpha, \Omega_\alpha)$ on X converging to (J, ω, Ω) and a family of embedded submanifolds $M_\alpha \subset X$ converging to $M_1 \cup M_2$ such that M_α is special Lagrangian in $(X, J_\alpha, \omega_\alpha, \Omega_\alpha)$.*

Proof. As mentioned earlier, the angle criterion is automatically satisfied when $n = 3$. It suffices to show that the homology condition in the Corollary implies the one in the Main Theorem. Since $[\Omega]$ spans $H^{3,0}(X)$, it follows that $[\text{Re } \Omega]$ and $[\text{Im } \Omega]$ span $H^{3,0}(X) \oplus H^{0,3}(X)$. By assumption, $\frac{[M_1]}{\text{Vol}(M_1)} - \frac{[M_2]}{\text{Vol}(M_2)} \neq 0$, and by the special Lagrangian condition, this real homology class is zero on both $[\text{Re } \Omega]$ and $[\text{Im } \Omega]$. So there exists some element of $\text{Re}[H^{2,1}(X) \oplus H^{1,2}(X)]$ which is not in the kernel of $\frac{[M_1]}{\text{Vol}(M_1)} - \frac{[M_2]}{\text{Vol}(M_2)}$. $\square$

Since a CY torus has trivial holonomy, our Main Theorem does not apply to this important case. However, we can still prove a version of the theorem in this setting.

Theorem 3 (Torus Version). *Let M_1 and M_2 be two embedded special Lagrangian submanifolds of a Calabi-Yau torus (T, J, ω, Ω) such that $\dim_{\mathbb{C}} T \geq 3$ and M_1 is flat. Assume that M_1 and M_2 intersect transversely at a single point p such that the tangent cone of $M_1 \cup M_2$ at p satisfies the angle criterion. Then there exists a family of Calabi-Yau structures $(J_\alpha, \omega, \Omega_\alpha)$ on T converging to (J, ω, Ω) and a family of embedded submanifolds $M_\alpha \subset T$ converging to $M_1 \cup M_2$ such that M_α is special Lagrangian in $(T, J_\alpha, \omega, \Omega_\alpha)$.*

The Main Theorem and the Torus Version share the hypothesis that M_1 and M_2 must intersect at a single point, but this condition is somewhat artificial. In light of the proof to follow, as long as there exists an *isolated* transverse intersection point p , we can still regularize the singularity at p , but the M_α 's will only be immersed rather than embedded. However, if $M_1 \cap M_2$ is a finite set of isolated transverse intersection points, all of which satisfy the angle criterion, then we can recover the embeddedness as follows: We first apply our result to one of these intersection points,¹ and then we apply Yng-Ing Lee's result on immersed SLags to each of the other intersection points. This procedure is possible because the property of being a transverse intersection point satisfying the angle criterion is an open condition with respect to the relevant topology. See Lemma 5.

It is a simple matter to construct infinitely many distinct pairs of flat SLag tori satisfying the angle criterion in the standard CY torus, $\mathbb{C}^n/\mathbb{Z}^{2n}$. Applying the discussion in the previous paragraph, we immediately obtain the following result.

Corollary 4. *There exist non-flat embedded special Lagrangian submanifolds of Calabi-Yau tori.*

With some extra work, the methods of this paper can probably be used to prove that our results hold in dimension two also. The methods of algebraic geometry should also apply in the dimension two case.

Concurrent with the writing of this paper, Joyce has produced some results on the general problem of desingularizing special Lagrangians with isolated conical singularities in almost Calabi-Yau manifolds [12, 13]. In particular, Theorem 7.11 of [13] combined with Lemma 16 of this paper and an understanding of the Lawlor necks can be used to prove the results of this paper. Note that Lemma 16 is the main ingredient of the Key Lemma of this paper. The methods used by Joyce are different from those presented here, and because of the added generality, the proofs are also more complicated.

ACKNOWLEDGEMENTS: I would like to thank Rick Schoen for suggesting the problem, listening to my ideas, and offering many helpful suggestions. This research was partially supported by a NSF Graduate Research Fellowship.

¹If either M_1 or M_2 is not connected, then we apply our result multiple times.

2 Preliminaries

The Main Theorem and the Torus Version share certain assumptions: We have two embedded special Lagrangian submanifolds M_1 and M_2 of a Calabi-Yau manifold (X, J, ω, Ω) with $\dim_{\mathbb{C}} X \geq 3$. We also assume that M_1 and M_2 intersect transversely at a single point p such that the tangent cone of $M_1 \cup M_2$ at p satisfies the angle criterion. This is the situation we assume from now until the proofs of the Key Lemma, which will depend on the additional assumptions in the two cases. We also assume without loss of generality that M_1 and M_2 are connected.

We now explain the idea behind these results. We wish to construct a family of approximate solutions M_α such that M_α converges to $M = M_1 \cup M_2$, M_α is exactly Lagrangian, and M_α is very close to being special Lagrangian. Once we have these M_α 's, we can construct small Hamiltonian deformations of them and hope that at least one of them is exactly special Lagrangian. This is actually too much to hope for, but we can add another degree of freedom to this deformation by simultaneously deforming the Calabi-Yau structure $(J_t, \omega_t, \Omega_t)$ and M_α itself so that the deformations of M_α remain Lagrangian with respect to ω_t . Using these deformations we define a deformation operator whose solutions correspond to special Lagrangians in $(X, J_t, \omega_t, \Omega_t)$. Using the Inverse Function Theorem together with some estimates, we obtain the desired solutions. The work lies in obtaining the appropriate estimates.

First we construct our family of approximate solutions M_α . This is where the angle criterion is relevant. Given $\phi_1, \dots, \phi_n \in \mathbb{R}$, we define $P(\phi_1, \dots, \phi_n)$ to be the oriented plane $[(\cos \phi_1) \frac{\partial}{\partial x^1} + (\sin \phi_1) \frac{\partial}{\partial y^1}] \wedge \dots \wedge [(\cos \phi_n) \frac{\partial}{\partial x^n} + (\sin \phi_n) \frac{\partial}{\partial y^n}]$. By the work of Lawlor [15], we know that for any $\theta_1, \dots, \theta_n \in (0, \pi)$ satisfying $\sum_{j=1}^n \theta_j = \pi$, there exists a special Lagrangian submanifold N of $\mathbb{C}^n$ that is asymptotic in an oriented sense to the two planes $P(0, \dots, 0)$ and $-P(-\theta_1, \dots, -\theta_n)$. This submanifold N has the property that ϵN converges to $[P(0, \dots, 0)] \cup [-P(-\theta_1, \dots, -\theta_n)]$ in an appropriate sense as $\epsilon \rightarrow 0$. These N 's, as well as their images under $SU(n) \times (\text{dilations})$ are called **Lawlor necks**. Note that there is also a Lawlor neck asymptotic in an oriented sense to the two planes $P(0, \dots, 0)$ and $-P(\theta_1, \dots, \theta_n)$.

Lemma 5. *If η and ξ are two special Lagrangian planes in $\mathbb{C}^n$, then there exists a Lawlor neck asymptotic in an oriented sense to η and ξ if and only if η and ξ are transverse planes satisfying the angle criterion. Moreover, both of these equivalent conditions are open conditions in the space of pairs of special Lagrangian planes. Finally, when $n \leq 3$, every pair of special Lagrangian planes satisfies the angle criterion.*

Proof. Let η and ξ be two transverse special Lagrangian planes in $\mathbb{C}^n$. The necessity of the angle criterion for the existence of a Lawlor neck is trivial, so we need only prove sufficiency. Assume that η and ξ satisfy the angle criterion. Without loss of generality, we may assume that $\eta = P(0, \dots, 0)$ by performing an $SU(n)$ change of coordinates. Since $-\xi$ is Lagrangian and transverse to η , we have $-\xi = P(\phi_1, \dots, \phi_n)$ for some $\phi_1, \dots, \phi_n \in (-\pi, 0) \cup (0, \pi)$, after an

$\mathrm{SO}(n)$ change of coordinates, where $\mathrm{SO}(n) \subset \mathrm{SU}(n)$ is the standard inclusion. See [2] for a proof of this fact. Note that we still have $\eta = P(0, \dots, 0)$ in this coordinate system. However, the unordered list of angles, $\phi_1, \dots, \phi_n$, is not uniquely determined. Adding or subtracting π to any of the ϕ_j 's merely changes the orientation, therefore doing this an even number of times leaves the oriented plane unchanged. We can find a canonical description by placing more restrictions on the angles. We demand that at most one of the ϕ_j 's has $|\phi_j| > \frac{\pi}{2}$, and if there is such a $|\phi_{j_0}|$, we demand that $|\phi_{j_0}| \leq \pi - |\phi_j|$ for all j . It is easy to verify that this prescription gives us a new unordered list of angles. Now observe that $|\phi_1|, \dots, |\phi_n|$ is precisely the unordered list of characterizing angles for the planes η and $-\xi$, therefore $\sum_{j=1}^n |\phi_j| = \pi$ by assumption. Since ξ is special Lagrangian, $\sum_{j=1}^n \phi_j \equiv \pi \pmod{2\pi}$. So we have $\sum_{j=1}^n \phi_j = \pm\pi$. In the positive case, we have $\sum_{j=1}^n (|\phi_j| - \phi_j) = 0$. Therefore all of the ϕ_j 's are positive and their sum is π . Similarly, in the negative case, we see that all of the ϕ_j 's are negative and their sum is $-\pi$. In either case, our brief discussion of Lawlor necks above shows that there exists a Lawlor neck asymptotic in an oriented sense to η and $-\xi$.

We now turn to the second statement in the Lemma. Let η and ξ be two transverse special Lagrangian planes satisfying the angle criterion. By the previous discussion, there exist $\mathrm{SU}(n)$ coordinates in which $\eta = P(0, \dots, 0)$ and $-\xi = P(\theta_1, \dots, \theta_n)$, where $\theta_1, \dots, \theta_n \in (0, \pi)$ and $\sum_{j=1}^n \theta_j = \pi$. (Or we have the negative case which is similar.) The key observation is that if ξ' is a SLag plane close to ξ , then we can make an $\mathrm{SO}(n) \subset \mathrm{SU}(n)$ change of coordinates so that $-\xi' = P(\theta'_1, \dots, \theta'_n)$, where $\theta'_1, \dots, \theta'_n$ is close to $\theta_1, \dots, \theta_n$ in the natural topology on unordered lists of n objects. This essentially follows from the fact that the unordered list of eigenvalues (with algebraic multiplicity) of a matrix continuously depends on the matrix. Since the change of coordinates was $\mathrm{SO}(n)$, we still have $\eta = P(0, \dots, 0)$ in this coordinate system. Since ξ' is special Lagrangian, $\sum_{j=1}^n \theta'_j \equiv \pi \pmod{2\pi}$, so for ξ' close enough to ξ , we see that $\sum_{j=1}^n \theta'_j = \pi$ and $\theta'_1, \dots, \theta'_n \in (0, \pi)$, thus proving the second statement of the Lemma.

The final statement in the Lemma about $n \leq 3$ is simple to verify. $\square$

Now we must use the existence of the local regularization to produce an approximate global regularization. The details of this construction are described in [1, 2, 16]. Here we only give a broad overview. Near the singular point p , we can choose a Darboux and normal coordinate system in a ball B around p such that $p = 0$ and Ω approaches dz appropriately as we approach 0. We know $M = M_1 \cup M_2$ becomes close to the tangent cone at p as we approach 0. The tangent cone must be a union of two planes calibrated by $\mathrm{Re} dz$. As long as this pair of planes meets the angle criterion, there exists a Lawlor neck N asymptotic to those two planes. For sufficiently small $\alpha > 0$ and certain constants C_δ and C_ϵ , choose

$$\delta = \frac{\alpha}{C_\delta} \text{ and } \epsilon = \frac{\alpha^{1+1/n}}{C_\epsilon}.$$

See [1, 2, 16] for the definitions of C_δ and C_ϵ .² These constants are chosen so that the following construction works and has the stated properties.

We can cut out a small ball $B_\delta(0)$ from M , glue a rescaled Lawlor neck ϵN into $B_\delta(0)$, and then interpolate in the annular region.³ This gives us M_α which we can think of as $M'_1 \cup T_1 \cup N' \cup T_2 \cup M'_2$, where $M'_i = M_i - B_\delta(0)$, N' is the rescaled Lawlor neck, and the T_i 's are the interpolated regions connecting M'_i to N' . It is evident that M_α converges to M by construction, and since each Lawlor neck has the topology of a cylinder, $S^{n-1} \times \mathbb{R}$, M_α is topologically the connected sum of M_1 and M_2 . We can choose the interpolation so that M_α is exactly Lagrangian. Since M_α is Lagrangian, it is a fact that at each point of M_α , $\Omega|_{M_\alpha} = e^{i\theta} \text{Vol}_{M_\alpha}$ for some θ . We call the multi-valued function θ the **Lagrangian angle function**. This “function” has the property that $J\nabla\theta$ is the mean curvature field H . On a special Lagrangian submanifold, $e^{i\theta} = 1$ and $H = 0$. One can show that M_α is approximately special Lagrangian in following sense [1, 2, 16].

Lemma 6. *Given the construction above, the Lagrangian angle function and the mean curvature field on M_α satisfy*

$$\begin{aligned} |\sin \theta|_0 + \alpha^\beta [\sin \theta]_\beta + \alpha |\nabla \sin \theta|_0 &\leq C\alpha \\ |1 - \cos \theta|_0 + \alpha^\beta [\cos \theta]_\beta + \alpha |\nabla \cos \theta|_0 &\leq C\alpha^2 \\ |H|_0 + \alpha^\beta [H]_\beta + \alpha |\nabla H|_0 &\leq C. \end{aligned}$$

Moreover, on $M_\alpha - B_\delta$, $e^{i\theta} = 1$ and $H = 0$.

3 The Deformation Operator

Now that we have our approximate solutions M_α , we can define the relevant deformation operator. Suppose that we have a smooth deformation $(J_t, \omega_t, \Omega_t)$ of the CY structure $(J, \omega, \Omega) = (J_0, \omega_0, \Omega_0)$ such that ω_t is always cohomologous to ω and $\langle [M_1] + [M_2], [\text{Im } \Omega_t] \rangle = 0$. We will choose the appropriate deformation of CY structure later in this paper; for now assume that we have already chosen it. By Moser's Theorem, there exists a smooth path of diffeomorphisms Ψ_t of X such that

$$\Psi_t^* \omega_t = \omega. \quad (2)$$

By the Lagrangian Neighborhood Theorem, let U be a tubular neighborhood of M_α symplectomorphic to T^*M_α so that we have a projection map $\pi : U \rightarrow M_\alpha$. Let τ be a smooth cutoff function supported in U such that $\tau = 1$ on $\frac{1}{2}U$, where $\frac{1}{2}U$ is defined using the structure of T^*M . Observe that we can choose U

²Throughout this paper α will be the parameter upon which most of our constructions depend. Because of this, we will explicitly write out the α dependence of all of our constants, with the only exceptions being δ and ϵ . We will use the letter C without subscript as a generic constant independent of α whose value may change even in a single chain of inequalities. For consistency we always use C as an upper bound.

³From now on we will write B_δ for $B_\delta(0)$ where there is no chance of confusion.

to have width greater than $\frac{\epsilon}{C}$ over $M_\alpha \cap B_{\epsilon r_0}$ for some r_0 and width greater than $\frac{1}{C}$ over $M'_1 \cup M'_2$, with an inverse linear interpolation in between. Now extend any function $h \in C^{2,\beta}(M_\alpha)$ to a function $\tilde{h} \in C^{2,\beta}(X)$ by defining $\tilde{h}(q) = \tau(q)h(\pi(q))$ on U and $\tilde{h} = 0$ outside U . Now define Φ_h to be the symplectomorphism generated by the Hamiltonian function $\tilde{h}$.

Definition. The **deformation operator** $F_\alpha : C^{2,\beta}(M_\alpha) \times \mathbb{R} \longrightarrow C^{0,\beta}(M_\alpha)$ is defined by

$$F_\alpha(h, t) = \langle (\Psi_t \circ \Phi_h)^*(\text{Im } \Omega_t), \text{Vol}_{M_\alpha} \rangle_{M_\alpha}$$

where the metric on M_α is the one induced by the Kähler metric on (X, J, ω) , independent of t .

Since M_α is a Lagrangian submanifold of (X, ω) and Φ_h is a symplectomorphism it follows that $\Phi_h(M_\alpha)$ is a Lagrangian submanifold of (X, ω) . Then by (2), it follows that $(\Psi_t \circ \Phi_h)(M_\alpha)$ is a Lagrangian submanifold of (X, ω_t) . Clearly, $F_\alpha(h, t) = 0$ iff $\text{Im } \Omega_t$ restricted to $(\Psi_t \circ \Phi_h)(M_\alpha)$ is identically zero. Therefore a solution of the equation $F_\alpha(h, t) = 0$ corresponds to a special Lagrangian submanifold of $(X, J_t, \omega_t, \Omega_t)$, and a small solution corresponds to a nearby special Lagrangian. So our goal is to show that for sufficiently small α , F_α has a small solution. Our method of constructing such solutions is the following version of the Inverse Function Theorem.

Theorem 7 (Inverse Function Theorem). *Let $F : \mathcal{B} \longrightarrow \mathcal{B}'$ be a C^1 map between Banach spaces and suppose that the linearization $DF(0)$ is an isomorphism. Moreover, assume that for some constants C_I , C_N , and r_1 , we have*

1. $\|DF(0)x\|_{\mathcal{B}'} \geq \frac{1}{C_I}\|x\|_{\mathcal{B}}$ for all $x \in \mathcal{B}$, and
2. $\|DF(0)x - DF(y)x\|_{\mathcal{B}'} \leq C_N\|x\|_{\mathcal{B}} \cdot \|y\|_{\mathcal{B}}$ for all $x, y \in \mathcal{B}$ with $\|y\|_{\mathcal{B}} < r_1$.

Then there exist neighborhoods U of 0 and V of $F(0)$ such that $F : U \longrightarrow V$ is a C^1 -diffeomorphism. Moreover, if $r \leq \min(r_1, (2C_I C_N)^{-1})$, then $B_{r/2C_I}(F(0)) \subset V$ and $B_{r/2C_I}(F(0)) \subset F(B_r(0))$.

In particular, when the hypotheses of the theorem are satisfied and additionally, $\|F(0)\|_{\mathcal{B}'} < \frac{r}{2C_I}$, we can solve the equation $F(y) = 0$ for some $\|y\|_{\mathcal{B}} < r$.

In order to invoke the Inverse Function Theorem in our situation, we need to choose our Banach spaces carefully. We define a smooth weight function ρ on M_α with the key property that the ball of radius $\rho(x)$ in M_α centered at x has uniformly bounded geometry. That is, in geodesic normal coordinates at x , we have $|g_{ij} - \delta_{ij}|_{1,\beta,B_{\rho(x)}(x)}^* \leq 1$ where the norm here is the local scale-invariant Schauder norm on $B_{\rho(x)}(x)$. We also require that $\tau = 1$ on the ball $B_{\rho(x)}(x, X)$. We can construct such a ρ with the following additional properties. See [1, 2, 16].

- For some r_0 and R independent of α ,

$$\rho(x) = \begin{cases} \epsilon R & \text{for } x \in N' = M_\alpha \cap B_{\epsilon r_0} \\ \text{interpolation} & \text{for } x \in M_\alpha \cap (B - B_{\epsilon r_0}) \\ R & \text{for } x \in M_\alpha - B \end{cases}$$

- $\rho(x) \leq C|x|$ for $x \in M_\alpha \cap (B - B_{\delta/2})$.
- $|\nabla \rho|_0 \leq C$.
- $\|\rho^{-1}\|_{L^2(M_\alpha)} \leq C$.

Definition. For any $0 < \beta < 1$, the ρ -weighted (k, β) -Schauder norm on $C^{k, \beta}(M_\alpha)$ is given by

$$|u|_{C_\rho^{k, \beta}(M_\alpha)} = |u|_{0, M_\alpha} + |\rho \nabla u|_{0, M_\alpha} + \cdots + |\rho^k \nabla^k u|_{0, M_\alpha} + [\rho^{k+\beta} \nabla^k u]_{\beta, M_\alpha}.$$

Let S be the first eigenfunction of the Laplacian on M_α , normalized so that $\|S\|_{L^2(M_\alpha)} = 1$.⁴ Then we define the Banach spaces $\mathcal{B}_{1, \alpha}$, $\mathcal{B}_\alpha$, and $\mathcal{B}'_\alpha$ as vector spaces

$$\begin{aligned} \mathcal{B}_{1, \alpha} &= \left\{ u \in C^{2, \beta}(M_\alpha) \left| \int_{M_\alpha} u = \int_{M_\alpha} u S = 0 \right. \right\} \\ \mathcal{B}_\alpha &= \mathcal{B}_{1, \alpha} \times \mathbb{R} \\ \mathcal{B}'_\alpha &= \left\{ u \in C^{0, \beta}(M_\alpha) \left| \int_{M_\alpha} u = 0 \right. \right\} \end{aligned}$$

with the norms

$$\begin{aligned} \|u\|_{\mathcal{B}_{1, \alpha}} &= |u|_{C_\rho^{2, \beta}(M_\alpha)} \\ \|(u, a)\|_{\mathcal{B}_\alpha} &= |u|_{C_\rho^{2, \beta}(M_\alpha)} + |a| \\ \|f\|_{\mathcal{B}'_\alpha} &= |\rho^2 f|_{C_\rho^{0, \beta}(M_\alpha)}. \end{aligned}$$

(The integrations above are taken with respect to the t -independent Kähler metric on M_α .)

Since Ψ_t and Φ_h are isotopies, $\langle [(\Psi_t \circ \Phi_h)(M_\alpha)], [\text{Im } \Omega_t] \rangle = \langle [M_\alpha], [\text{Im } \Omega_t] \rangle = \langle [M_1] + [M_2], [\text{Im } \Omega_t] \rangle = 0$, and therefore $F_\alpha(\mathcal{B}_\alpha) \subset \mathcal{B}'_\alpha$. From now on we think of the **deformation operator** F_α as an operator from $\mathcal{B}_\alpha$ to $\mathcal{B}'_\alpha$.

The choice of β is not particularly important; it is simply a small constant independent of α . The purpose of the weighted norm is to achieve estimates that scale nicely with respect to α . The reason why we take the orthogonal complement of the functions 1 and S is that 1 lies in the kernel of the linearization, and S lies in the approximate kernel of the linearization.

Let us summarize what we need to prove in order to invoke the Inverse Function Theorem argument:

- We need an injectivity estimate on $DF_\alpha(0, 0)$; we must establish the existence of a constant $C_I(\alpha)$ as in condition 1 of the Inverse Function Theorem and find its dependence on α .
- We need to show that $DF_\alpha(0, 0)$ is surjective.

⁴In contrast to our use of constants, many geometric objects such as functions and operators will depend on α , but we will suppress this dependence in the notation for the purpose of readability. The loss of clarity should be minimal since these objects are all defined on M_α .

- We need a nonlinear estimate; we must establish the existence of a constant $C_N(\alpha)$ as in condition 2 of the Inverse Function Theorem and find its dependence on α .
- We need to bound $F_\alpha(0, 0)$ in terms of α .

We first compute $DF_\alpha(0, 0)$.

Proposition 8.

$$DF_\alpha(0, 0)(u, a) = \Delta u + Pu + a\psi \quad (3)$$

where $P : \mathcal{B}_{1,\alpha} \longrightarrow \mathcal{B}'_\alpha$ is given by

$$Pu = (\cos \theta - 1)\Delta u - (\sin \theta)\langle H, J\nabla u \rangle$$

and $\psi \in \mathcal{B}'_\alpha$ is given by

$$\psi = \langle L_V(\text{Im } \Omega) + \text{Im } \dot{\Omega}, \text{Vol}_{M_\alpha} \rangle$$

where V is the vector field generating the flow Ψ_t at time $t = 0$, and $\dot{\Omega} = \frac{d}{dt}\Omega_t|_{t=0}$.

The calculation of ψ is self-evident. The rest of the calculation is straightforward and can be found in [1] and [16]. The reason we write Δ and P separately in equation (3) is that the P term turns out to be negligible, and therefore it suffices to understand Δ and ψ . The unimportance of P is expressed in the following lemma.

Lemma 9. For sufficiently small α , for any $u \in \mathcal{B}_{1,\alpha}$,

$$\|Pu\|_{\mathcal{B}'_\alpha} \leq C\alpha^{1-\beta}\|u\|_{\mathcal{B}_{1,\alpha}}.$$

Proof. The proof essentially follows directly from the bounds given in Lemma 6.

$$\begin{aligned} |\rho^2(1 - \cos \theta)\Delta u|_0 &\leq |1 - \cos \theta|_0 \cdot |\rho^2 \Delta u|_0 \\ &\leq C\alpha^2 |u|_{C_\rho^{2,\beta}}. \end{aligned}$$

$$\begin{aligned} [\rho^{2+\beta}(1 - \cos \theta)\Delta u]_\beta &\leq [1 - \cos \theta]_\beta \cdot |\rho^{2+\beta} \Delta u|_0 + |1 - \cos \theta|_0 \cdot [\rho^{2+\beta} \Delta u]_\beta \\ &\leq (C\alpha^{2-\beta}R^\beta + C\alpha^2)|u|_{C_\rho^{2,\beta}}. \end{aligned}$$

$$\begin{aligned} |\rho^2(\sin \theta)\langle H, J\nabla u \rangle|_0 &\leq |\rho(\sin \theta)H|_0 \cdot |\rho J\nabla u|_0 \\ &\leq RC\alpha |u|_{C_\rho^{2,\beta}}. \end{aligned}$$

For the final inequality, we use the fact that $[v]_\beta \leq |v|_0 + |\nabla v|_0$.

$$\begin{aligned} [\rho^{2+\beta}(\sin \theta)\langle H, J\nabla u \rangle]_\beta &\leq [\sin \theta]_\beta \cdot |H|_0 \cdot |\rho^{2+\beta} J\nabla u|_0 + |\sin \theta|_0 \cdot [H]_\beta \cdot |\rho^{2+\beta} J\nabla u|_0 + \\ &\quad |\sin \theta|_0 \cdot |H|_0 \cdot [\rho^{2+\beta} J\nabla u]_\beta \\ &\leq (C\alpha^{1-\beta})CR^{1+\beta}|\rho \nabla u|_0 + (C\alpha)(C\alpha^{-\beta})R^{1+\beta}|\rho \nabla u|_0 + \\ &\quad (C\alpha)C(|\rho^{2+\beta} \nabla u|_0 + |(\nabla \rho^{2+\beta})(\nabla u)|_0 + |\rho^{2+\beta} \nabla^2 u|_0) \\ &\leq C\alpha^{1-\beta}|u|_{C_\rho^{2,\beta}} + C\alpha(R^{1+\beta} + CR^\beta + R^\beta)|u|_{C_\rho^{2,\beta}} \end{aligned}$$

where the last line uses the bound on $|\nabla \rho|_0$. Now combine the previous four inequalities to deduce the desired result. $\square$

4 Analysis of the Laplacian on M_α

The first step in establishing an injectivity estimate for the linearized deformation operator is finding a lower bound for the second eigenvalue of the Laplacian. The second step is to combine this lower bound with an elliptic estimate to obtain an injectivity estimate for $\Delta : \mathcal{B}_{1,\alpha} \rightarrow \mathcal{B}'_\alpha$.

It is a fact that on any Riemannian manifold M , for any $f \in L^2(M)$ with one derivative in $L^2(M)$ such that $\int_M f = 0$, we have $\int_M |\nabla f|^2 \geq \lambda_1(M) \int_M f^2$, where $\lambda_1(M)$ is the first eigenvalue of the Laplacian. From this it follows easily that if we drop the condition $\int_M f = 0$, then we have

$$\lambda_1(M) \leq \frac{\int_M |\nabla f|^2}{\int_M f^2 - \frac{1}{\text{Vol}(M)} (\int_M f)^2}. \quad (4)$$

We define a smooth cutoff function φ on M_α with the following properties: $\varphi = 0$ in B_δ , $\varphi = 1$ outside $B_{2\delta}$, and $|\nabla \varphi| \leq \frac{C}{\delta}$ for some C . Recall that $M_\alpha - B_\delta = (M_1 \cup M_2) - B_\delta$, and therefore we may think of φ as a function on either M_α or on $M_1 \amalg M_2$. Observe that because we have uniformly bounded mean curvature, the Monotonicity Formula provides the following bounds which we will use repeatedly:

$$\begin{aligned} \text{Vol}(M_\alpha \cap B_{2\delta}) &= O(\delta^n) \\ \text{Vol}(M_i \cap B_{2\delta}) &= O(\delta^n) \\ \text{Vol}(M_i - B_{2\delta}) &\leq C \\ \text{Vol}(M_i - B_{2\delta}) &\geq \frac{1}{C} \end{aligned}$$

Lemma 10. *For small enough α ,*

$$\lambda_1(M_\alpha) \leq C\delta^{n-2}.$$

Proof. Let

$$f = \varphi \left[\frac{\mathcal{X}_{M_1 - B_\delta}}{\text{Vol}(M_1 - B_{2\delta})} - \frac{\mathcal{X}_{M_2 - B_\delta}}{\text{Vol}(M_2 - B_{2\delta})} \right]$$

where $\mathcal{X}_A$ denotes the characteristic function of A . Clearly, $\int_{M_\alpha} f = O(\delta^n)$. Using inequality (4), we see that

$$\begin{aligned} \lambda_1(M_\alpha) &\leq \frac{\int_{M_\alpha} |\nabla f|^2}{\int_{M_\alpha} f^2 - \frac{1}{\text{Vol}(M_\alpha)} (\int_{M_\alpha} f)^2} \\ &\leq \frac{C \int_{M_\alpha \cap B_{2\delta}} \delta^{-2}}{\frac{1}{\text{Vol}(M_1 - B_{2\delta})} + \frac{1}{\text{Vol}(M_2 - B_{2\delta})} - O(\delta^{2n})} \text{ by the properties of } \varphi \\ &\leq C\delta^{n-2} \text{ since the denominator is bounded below.} \end{aligned}$$

□

We would like to have some idea of what S looks like. By the previous Lemma together with Lemma 5 of Yng-Ing Lee's paper [16], we know that $|S|_0$ is bounded independently of α .⁵ This fact allows us to use our knowledge of the kernel of the Laplacian on $M_1 \amalg M_2$ to construct a function that approximates S in the L^2 sense.

Lemma 11. *Define*

$$\bar{S} = a_1 \mathcal{X}_{M_1} + a_2 \mathcal{X}_{M_2}$$

where

$$a_1 = \frac{1}{\text{Vol}(M_1)} \sqrt{\frac{\text{Vol}(M_1) \text{Vol}(M_2)}{\text{Vol}(M_1) + \text{Vol}(M_2)}} \text{ and } a_2 = \frac{-1}{\text{Vol}(M_2)} \sqrt{\frac{\text{Vol}(M_1) \text{Vol}(M_2)}{\text{Vol}(M_1) + \text{Vol}(M_2)}}.$$

Then for small enough α ,

$$\|S - \varphi \bar{S}\|_{L^2(M_\alpha)} \leq C \delta^{(n-2)/2}$$

and

$$\|\bar{S} - \varphi \bar{S}\|_{L^2(M_1 \amalg M_2)} \leq C \delta^{n/2}.$$

Proof. First, the bound on S implies that

$$\|S - \varphi S\|_{L^2(M_\alpha)}^2 \leq \|S\|_{L^2(M_\alpha \cap B_{2\delta})}^2 = O(\delta^n). \quad (5)$$

Since φS is defined on $M_1 \amalg M_2$, we have

$$\begin{aligned} \int_{M_1 \amalg M_2} |\nabla(\varphi S)|^2 &= \int_{M_\alpha} |\nabla(\varphi S)|^2 \\ &\leq 2 \int_{M_\alpha \cap B_{2\delta}} |\nabla \varphi|^2 S^2 + 2 \int_{M_\alpha} \varphi^2 |\nabla S|^2 \\ &\leq O(\delta^{n-2}) + 2 \int_{M_\alpha} |\nabla S|^2 \text{ arguing as in Lemma 10} \\ &= O(\delta^{n-2}) \text{ by Lemma 10 and the normalization of } S. \end{aligned}$$

Note that $M_1 \amalg M_2$ has a two-dimensional kernel spanned by $\mathcal{X}_{M_1}$ and $\mathcal{X}_{M_2}$, and its first non-zero eigenvalue is obviously a constant independent of α . Therefore the estimate above shows that if $a'_1 \mathcal{X}_{M_1} + a'_2 \mathcal{X}_{M_2}$ is the orthogonal projection of φS onto the kernel, then

$$\begin{aligned} \|\varphi S - (a'_1 \mathcal{X}_{M_1} + a'_2 \mathcal{X}_{M_2})\|_{L^2(M_1 \amalg M_2)}^2 &\leq C \|\nabla[\varphi S - (a'_1 \mathcal{X}_{M_1} + a'_2 \mathcal{X}_{M_2})]\|_{L^2(M_1 \amalg M_2)}^2 \\ &= C \|\nabla(\varphi S)\|_{L^2(M_1 \amalg M_2)}^2 \\ &= O(\delta^{n-2}) \text{ by the previous calculation.} \end{aligned} \quad (6)$$

The bounds (5) and (6) show that

$$\frac{(a'_1)^2 \text{Vol}(M_1) + (a'_2)^2 \text{Vol}(M_2)}{\|a'_1 \mathcal{X}_{M_1} + a'_2 \mathcal{X}_{M_2}\|_{L^2(M_1 \amalg M_2)}^2} = 1 + O(\delta^{n-2})$$

⁵This Lemma depends on the Michael-Simon Inequality and uses the fact that the mean curvature of M_α is bounded independently of α .

and

$$a'_1 \text{Vol}(M_1) + a'_2 \text{Vol}(M_2) = \int_{M_1 \amalg M_2} a'_1 \mathcal{X}_{M_1} + a'_2 \mathcal{X}_{M_2} = O(\delta^{(n-2)/2}).$$

Solving these equations, we find that

$$a'_1 = a_1 + O(\delta^{(n-2)/2}) \text{ and } a'_2 = a_2 + O(\delta^{(n-2)/2})$$

where a_1 and a_2 were defined above.⁶ It now follows that

$$\|(a'_1 \mathcal{X}_{M_1} + a'_2 \mathcal{X}_{M_2}) - \bar{S}\|_{L^2(M_1 \amalg M_2)}^2 = O(\delta^{n-2}). \quad (7)$$

Finally, similar to the bound (5), we see that

$$\|\bar{S} - \varphi \bar{S}\|_{L^2(M_1 \amalg M_2)}^2 = O(\delta^n). \quad (8)$$

Putting together the bounds (5), (6), (7), and (8), we obtain the desired result. $\square$

We can now use our knowledge of S to help us show that the second eigenvalue of the Laplacian on M_α is bounded below.

Proposition 12 (Second Eigenvalue Estimate). *For small enough α , the second eigenvalue of the Laplacian of M_α , $\lambda_2(M_\alpha)$, is bounded below. In particular, for each $u \in \mathcal{B}_{1,\alpha}$,*

$$\|\Delta u\|_{L^2(M_\alpha)} \geq \frac{1}{C} \|u\|_{L^2(M_\alpha)}$$

Proof. Let f be an eigenfunction for the second eigenvalue of the Laplacian, normalized so that $\|f\|_{L^2(M_\alpha)} = 1$. Using the min-max characterization of $\lambda_2(M_\alpha)$ one can show that $\lambda_2(M_\alpha)$ is bounded above independently of α . This fact allows us to apply Lemma 5 of Yng-Ing Lee's paper [16] to show that $|f|_0$ is bounded independently of α . We compute

$$\begin{aligned} \lambda_2(M_\alpha) &= \int_{M_\alpha} |\nabla f|^2 \\ &= \int_{M_\alpha} |\nabla(\varphi f) + \nabla((1-\varphi)f)|^2 \\ &\geq \int_{M_1} |\nabla(\varphi f)|^2 + \int_{M_2} |\nabla(\varphi f)|^2 - 2 \int_{M_\alpha \cap B_{2\delta}} |\nabla(1-\varphi)|^2 f^2 - 2 \int_{M_\alpha} (1-\varphi)^2 |\nabla f|^2 \\ &\geq \int_{M_1} |\nabla(\varphi f)|^2 + \int_{M_2} |\nabla(\varphi f)|^2 - O(\delta^{n-2}) - 2 \int_{M_\alpha} |\nabla f|^2. \end{aligned}$$

⁶Of course, the solution is only determined up to a sign, but there was a sign ambiguity in our original definition of S , so we can simply define S to have the sign consistent with these equations.

Therefore,

$$\begin{aligned}
3\lambda_2(M_\alpha) &\geq \int_{M_1} |\nabla(\varphi f)|^2 + \int_{M_2} |\nabla(\varphi f)|^2 - O(\delta^{n-2}) \\
&\geq \lambda_1(M_1) \left[\int_{M_1} \varphi^2 f^2 - \frac{1}{\text{Vol}(M_1)} \left(\int_{M_1} \varphi f \right)^2 \right] + \\
&\quad \lambda_2(M_2) \left[\int_{M_2} \varphi^2 f^2 - \frac{1}{\text{Vol}(M_2)} \left(\int_{M_2} \varphi f \right)^2 \right] - O(\delta^{n-2}) \text{ by inequality (4)} \\
&\geq C \int_{M_\alpha} f^2 - O(\delta^n) - \frac{\lambda_1(M_1)}{\text{Vol}(M_1)} \left(\int_{M_1} \varphi f \right)^2 \\
&\quad - \frac{\lambda_2(M_2)}{\text{Vol}(M_2)} \left(\int_{M_2} \varphi f \right)^2 - O(\delta^{n-2}) \\
&\geq C - C \left[\left(\int_{M_1} \varphi f \right)^2 + \left(\int_{M_2} \varphi f \right)^2 \right] - O(\delta^{n-2}).
\end{aligned}$$

Now we must bound the terms in brackets in the previous line.

$$\begin{aligned}
\left| \int_{M_1} \varphi f \right| &= \left| \int_{M_1 \amalg M_2} \left(\frac{\bar{S} - a_2}{a_1 - a_2} \right) \varphi f \right| \text{ by the definition of } \bar{S} \text{ from Lemma 11} \\
&= \frac{1}{a_1 - a_2} \left| \int_{M_\alpha} (\varphi \bar{S}) f - a_2 \int_{M_\alpha} \varphi f \right| \\
&= \frac{1}{a_1 - a_2} \left| \int_{M_\alpha} S f + \int_{M_\alpha} (\varphi \bar{S} - S) f - a_2 \int_{M_\alpha} f + O(\delta^n) \right| \\
&\leq C \|\varphi \bar{S} - S\|_{L^2(M_\alpha)} + O(\delta^n) \text{ since } \int_{M_\alpha} S f = \int_{M_\alpha} f = 0 \\
&= O(\delta^{(n-2)/2}) + O(\delta^n) \text{ by Lemma 11.}
\end{aligned}$$

The estimate for M_2 is similar. $\square$

Recall that the weight function ρ was chosen so that M_α has uniformly bounded geometry in $\rho(x)$ neighborhoods of x in M_α . This allows us to use the local scale-invariant elliptic Schauder estimate to deduce a global elliptic Schauder estimate independent of α . We omit the proof which is standard and straightforward.

Proposition 13 (Global Elliptic Schauder Estimate). *For sufficiently small α , for any $u \in C^{2,\beta}(M_\alpha)$,*

$$|u|_{C_\rho^{2,\beta}} \leq C(|\rho^2 \Delta u|_{C_\rho^{0,\beta}} + |u|_0).$$

The following lemma, proved in [16], translates the second eigenvalue estimate from the L^2 setting to the Schauder setting.⁷ Choose any $\nu > 0$ independent of α .

⁷The proof of this lemma follows easily from a De Giorgi-Nash estimate, which in turn depends on the Michael-Simon inequality and bounded mean curvature.

Lemma 14. *For sufficiently small α ,*

$$|u|_0 \leq \epsilon^{-\nu} |\rho^2 \Delta u|_{C_\rho^{0,\beta}}$$

for all $u \in \mathcal{B}_{1,\alpha}$.

Combining this Lemma with Global Elliptic Estimate immediately leads us to an injectivity estimate for $\Delta : \mathcal{B}_{1,\alpha} \longrightarrow \mathcal{B}'_\alpha$.

Proposition 15 (Laplacian Injectivity Estimate). *For sufficiently small α , for all $u \in \mathcal{B}_{1,\alpha}$,*

$$|u|_{C_\rho^{2,\beta}} \leq C \epsilon^{-\nu} |\rho^2 \Delta u|_{C_\rho^{0,\beta}}.$$

5 Proofs of the Key Lemma

It is well-known that Δ is an isomorphism from $\mathcal{B}_{1,\alpha} \oplus \langle S \rangle$ to $\mathcal{B}'_\alpha$, but because of the small first eigenvalue, we had to remove S in order to obtain a good injectivity estimate. Of course, removing S costs us surjectivity. We added the extra degree of freedom in order to restore surjectivity. The essential requirement of the extra degree of freedom is that its linearization ψ must have a significant S component. This is the content of our Key Lemma. We now construct a deformation $(J_t, \omega_t, \Omega_t)$ so that ψ has the desired property.

Lemma 16. *Under the assumptions of the Main Theorem or the Torus Version, there exists a deformation of Calabi-Yau structure $(J_t, \omega_t, \Omega_t)$ such that ω_t is cohomologous to ω , $\langle [M_1] + [M_2], [\text{Im } \Omega_t] \rangle = 0$, and*

$$\left\langle \frac{[M_1]}{\text{Vol}(M_1)} - \frac{[M_2]}{\text{Vol}(M_2)}, [\text{Im } \dot{\Omega}] \right\rangle \neq 0$$

for sufficiently small α .

Proof of Main Case. By a result of Tian [20] and Todorov [21], the first-order deformation space of complex structures on (X, J) is exactly the space of harmonic $(n-1, 1)$ forms, $\mathcal{H}^{n-1,1}(X)$, and all of these first-order deformations extend to actual deformations. By the hypotheses of the Main Theorem, we can find $\lambda \in \text{Re}[\mathcal{H}^{n-1,1}(X) \oplus \mathcal{H}^{1,n-1}(X)]$ such that $\left\langle \frac{[M_1]}{\text{Vol}(M_1)} - \frac{[M_2]}{\text{Vol}(M_2)}, [\lambda] \right\rangle \neq 0$. Since $\mathcal{H}^{n-1,1}(X)$ and $\mathcal{H}^{1,n-1}(X)$ are complex conjugate to each other, we can certainly find $\chi \in \mathcal{H}^{n-1,1}(X)$ such that $\text{Im } \chi = \lambda$. Choose a complex structure deformation J_t whose first-order deformation is χ . Recall that the choice of J_t determines the holomorphic $(n, 0)$ -form Ω_t up to a constant. It is a fact that the effect of the first-order deformation χ on $\dot{\Omega}$ is expressed by the formula

$$\dot{\Omega} = c\Omega + \chi.$$

See Candelas and de la Ossa [3] for details. Since $\text{Im } \Omega$ vanishes on M_1 and M_2 , and $\text{Re } \Omega$ calibrates M_1 and M_2 , it follows that $\left\langle \frac{[M_1]}{\text{Vol}(M_1)} - \frac{[M_2]}{\text{Vol}(M_2)}, [\Omega] \right\rangle = 0$, and therefore $\left\langle \frac{[M_1]}{\text{Vol}(M_1)} - \frac{[M_2]}{\text{Vol}(M_2)}, [\text{Im } \dot{\Omega}] \right\rangle \neq 0$.

We know that $[\omega]$ lies in the Kähler cone of (X, J) . Since (X, J) admits a Calabi-Yau structure with holonomy equal to $SU(n)$, so does (X, J_t) . In this case, it is a fact that $H^{2,0}(X, J_t) = H^{0,2}(X, J_t) = 0$. See Joyce [14] for details. Therefore the Kähler cone of (X, J_t) is open in $H^2(X)$ for each t . So for small t , $[\omega]$ lies in the Kähler cone of (X, J_t) . Therefore we can find a unique $\omega_t \in [\omega]$ such that (X, J_t, ω_t) admits a CY structure. This determines Ω_t up to a phase, and we choose the phase so that $\langle [M_1] + [M_2], [\text{Im } \Omega_t] \rangle = 0$. $\square$

Before we discuss the Torus Case, let us study Calabi-Yau tori and the moduli space of CY tori.

Proposition 17. *We can characterize all Calabi-Yau tori and all flat special Lagrangian submanifolds of them as follows.*

1. *Given a rank $2n$ lattice $\Gamma \subset \mathbb{C}^n$, we define a Calabi-Yau structure on $\mathbb{C}^n/\Gamma$ using the standard Calabi-Yau structure on $\mathbb{C}^n$. Every Calabi-Yau torus T is given by this construction.*
2. *Given a special Lagrangian plane η invariant with respect to some rank n sublattice of Γ , we obtain a special Lagrangian torus η/Γ in $\mathbb{C}^n/\Gamma$. Every flat special Lagrangian submanifold of T is a union of special Lagrangian tori given by this construction.*

Proof. Let (T, J, ω, Ω) be a CY torus. Clearly, we have an induced CY structure on the universal cover $(\mathbb{R}^{2n}, \tilde{J}, \tilde{\omega}, \tilde{\Omega})$. Since the Kähler metric on T has zero Ricci curvature, we can apply Cheeger and Gromoll's theorem on manifolds with non-negative Ricci curvature [4] to see that the induced metric on the universal cover is flat. Choose a point $q \in \mathbb{R}^{2n}$. We can find a basis $e_1, \dots, e_{2n}$ of $T_q \mathbb{R}^{2n}$ such that if $e^1, \dots, e^{2n}$ is the dual basis, then $\tilde{J}e_j = e_{j+n}$ for $j \leq n$, $\tilde{J}e_j = -e_{j-n}$ for $j > n$, $\tilde{\omega} = \sum_{j=1}^n e^j \wedge e^{j+n}$, and $\tilde{\Omega} = (e_1 + ie_{n+1}) \wedge \dots \wedge (e_n + ie_{2n})$ at the point q . Since the metric is flat, we can extend $e_1, \dots, e_{2n}$ to a frame field over all of $\mathbb{R}^{2n}$ in a unique way via parallel translation. Note that $[e_j, e_k] = 0$ everywhere, so we can find a global coordinate system $x^1, \dots, x^n, y^1, \dots, y^n$ such that $\frac{\partial}{\partial x^j} = e_j$ and $\frac{\partial}{\partial y^j} = e_{j+n}$. Since $\nabla J = \nabla \omega = \nabla \Omega = 0$, it follows that $\tilde{J}(\frac{\partial}{\partial x^j}) = \frac{\partial}{\partial y^j}$, $\tilde{J}(\frac{\partial}{\partial y^j}) = -\frac{\partial}{\partial x^j}$, $\tilde{\omega} = \sum_{j=1}^n dx^j \wedge dy^j$, and $\tilde{\Omega} = dz^1 \wedge \dots \wedge dz^n$ everywhere. That is, the induced CY structure on the universal cover is the standard CY structure on $\mathbb{C}^n$. Moreover, since each Deck transformation must be an orientation-preserving isometry, the Deck transformations are translations. $\square$

Let M be a flat SLAG in T . Since M is flat and minimal in T , which is also flat, it is a simple consequence of the Gauss equation that M is totally geodesic in T . In fact, Ricci-flatness would have been sufficient. See [5]. Therefore, locally, the lift of M up to $\mathbb{C}^n$ is a piece of a SLAG plane. It follows that M is the quotient of a union of SLAG planes which are invariant under some rank n sublattice of Γ . $\square$

We know that the CY structure of a torus is determined by a lattice. The space of all rank $2n$ lattices in $\mathbb{C}^n$ is $GL(2n, \mathbb{R})/SL(2n, \mathbb{Z})$ where $SL(2n, \mathbb{Z})$ acts

on the right. Since the group of CY structure preserving automorphisms of $\mathbb{C}^n$ is $SU(n) \ltimes (\text{translations})$, it follows that the global moduli space of Calabi-Yau tori is precisely $GL(2n, \mathbb{R})/SL(2n, \mathbb{Z})$ modulo the action of $SU(n)$ on the left. Therefore, locally, the deformation space is simply a neighborhood of the identity in $GL(2n, \mathbb{R})/SU(n)$ where $SU(n)$ acts on the left. Ignoring the (discrete) redundancies arising from the lattice automorphisms, $SL(2n, \mathbb{Z})$, the space of possible complex structures corresponds to $GL(2n, \mathbb{R})/GL(n, \mathbb{C})$, and after this choice is made, the space of compatible symplectic structures corresponds to $GL(n, \mathbb{C})/U(n)$. On the other hand, we can choose the symplectic structure from the space $GL(2n, \mathbb{R})/Sp(2n, \mathbb{R})$ and then choose a compatible complex structure from the space $Sp(2n, \mathbb{R})/U(n)$. Finally, of course, we choose a phase for Ω from $U(n)/SU(n)$. Finally, observe that changing the lattice by $A \in GL(2n, \mathbb{R})$ has the same effect as keeping the lattice and the canonical local coordinate systems on T fixed, but changing the CY structure to $(A^{-1}JA, A^*\omega, A^*\Omega)$ with respect to those coordinates. This is the point of view we adopt in the following proof, and we will no longer mention lattices.

Proof of Torus Case of Lemma 16. By Proposition 17, any connected, flat SLAG M_1 in (T, J, ω, Ω) is actually a SLAG torus with constant tangent plane with respect to the canonical local coordinates, so we can perform an $SU(n)$ change of coordinates taking M_1 to a SLAG torus with tangent plane $\frac{\partial}{\partial x^1} \wedge \dots \wedge \frac{\partial}{\partial x^n}$ at each point of M_1 . Since the $SU(n)$ change of coordinates preserves the CY structure, we may assume without loss of generality that $T_q M_1 = \frac{\partial}{\partial x^1} \wedge \dots \wedge \frac{\partial}{\partial x^n}$ at each $q \in M_1$.

Consider a deformation $A_t^{-1}JA_t$ of the complex structure. Then the holomorphic $(n, 0)$ form is $A_t^*\Omega$ up to a constant. A simple calculation then shows that $\text{Im } \tilde{\Omega} = c\Omega + \chi$, where

$$\chi = \sum_{j,k=1}^n B_{jk} dz^1 \wedge \dots \wedge \overbrace{d\bar{z}^j}^{k\text{-th spot}} \wedge \dots \wedge dz^n$$

and $B_{jk} = \dot{A}_{k\bar{j}}$. Another simple calculation shows that if $A_t \in Sp(2n, \mathbb{R})$, then the corresponding B must be (complex) symmetric, and conversely, for any (complex) symmetric B , we can find $A_t \in Sp(2n, \mathbb{R})$ such that $B_{jk} = \dot{A}_{k\bar{j}}$.

In particular, we can find $A_t \in Sp(2n, \mathbb{R})$ and a phase $\phi_t \in \mathbb{R}$ such that $(A_t^{-1}JA_t, \omega, \Omega_t = e^{i\phi_t} A_t^*\Omega)$ is a CY structure with $\langle [M_1] + [M_2], [\text{Im } \Omega_t] \rangle = 0$, and

$$\chi = i \sum_{j=1}^n dz^1 \wedge \dots \wedge \overbrace{d\bar{z}^j}^{j\text{-th spot}} \wedge \dots \wedge dz^n.$$

Note that

$$\text{Im } \chi = \sum_{j=1}^n \text{Re}(dz^1 \wedge \dots \wedge \overbrace{d\bar{z}^j}^{j\text{-th spot}} \wedge \dots \wedge dz^n).$$

It remains to show that $\left\langle \frac{[M_1]}{\text{Vol}(M_1)} - \frac{[M_2]}{\text{Vol}(M_2)}, [\text{Im } \chi] \right\rangle \neq 0$. First, observe that $\text{Re}(dz^1 \wedge \dots \wedge d\bar{z}^j \wedge \dots \wedge dz^n)$ is a calibration for each j , and that $\frac{\partial}{\partial x^1} \wedge \dots \wedge \frac{\partial}{\partial x^n}$ is calibrated by each of these calibrations. Therefore $\frac{1}{\text{Vol}(M_1)} \int_{M_1} \text{Im } \chi = n$ and $\frac{1}{\text{Vol}(M_2)} \int_{M_2} \text{Im } \chi \leq n$ with equality iff M_2 is also calibrated by each of the calibrations $\text{Re}(dz^1 \wedge \dots \wedge d\bar{z}^j \wedge \dots \wedge dz^n)$. We momentarily consider the “torus” planes ξ of the form

$$\left[\left[(\cos \theta_1) \frac{\partial}{\partial x^1} + (\sin \theta_1) \frac{\partial}{\partial y^1} \right] \wedge \dots \wedge \left[(\cos \theta_n) \frac{\partial}{\partial x^n} + (\sin \theta_n) \frac{\partial}{\partial y^n} \right] \right].$$

It is easy to verify that if ξ is calibrated by $\text{Re } \Omega = \text{Re}(dz^1 \wedge \dots \wedge dz^n)$ and $\text{Re}(dz^1 \wedge \dots \wedge d\bar{z}^j \wedge \dots \wedge dz^n)$ for each j , then $\xi = \frac{\partial}{\partial x^1} \wedge \dots \wedge \frac{\partial}{\partial x^n}$. Then by Morgan’s Torus Lemma, it follows that $\frac{\partial}{\partial x^1} \wedge \dots \wedge \frac{\partial}{\partial x^n}$ is the only plane simultaneously calibrated by $\text{Re}(dz^1 \wedge \dots \wedge dz^n)$ and $\text{Re}(dz^1 \wedge \dots \wedge d\bar{z}^j \wedge \dots \wedge dz^n)$ for each j . See [6] for more on Morgan’s Torus Lemma. Therefore if M_2 is calibrated by $\text{Re } \Omega = \text{Re}(dz^1 \wedge \dots \wedge dz^n)$ and $\text{Re}(dz^1 \wedge \dots \wedge d\bar{z}^j \wedge \dots \wedge dz^n)$ for each j , it follows that $T_p M_2 = \frac{\partial}{\partial x^1} \wedge \dots \wedge \frac{\partial}{\partial x^n}$, violating the transversality assumption. $\square$

Lemma 18 (Key Lemma). *For small α ,*

$$\left| \int_{M_\alpha} \psi S \right| \geq \frac{1}{C}. \quad (9)$$

Proof. Recall that $\psi = \langle L_V(\text{Im } \Omega), \text{Vol}_{M_\alpha} \rangle + \langle \text{Im } \dot{\Omega}, \text{Vol}_{M_\alpha} \rangle$. We’ll show that the first term integrated against S is small while the second term integrated against S is bounded below. Since Ω is closed, $L_V(\text{Im } \Omega) = d(V \lrcorner \text{Im } \Omega)$. Note that the two differential forms $V \lrcorner \text{Im } \Omega$ and $d(V \lrcorner \text{Im } \Omega)$ are defined on X independently of α . From this it follows that $|V \lrcorner \text{Im } \Omega|_{M_\alpha}|_0$ and $|d(V \lrcorner \text{Im } \Omega)|_{M_\alpha}|_0$ are bounded independently of α , where these are the induced norms on M_α . Thus

$$\begin{aligned} \left| \int_{M_\alpha} (d(V \lrcorner \text{Im } \Omega)|_{M_\alpha}) S \right| &\leq \left| \int_{M_\alpha} (d(V \lrcorner \text{Im } \Omega)|_{M_\alpha}) \varphi \bar{S} \right| + C\delta^{(n-2)/2} \text{ by Lemma 11} \\ &= \left| \int_{M_\alpha} (V \lrcorner \text{Im } \Omega) d(\varphi \bar{S}) \right| + C\delta^{(n-2)/2} \\ &\leq \left| \int_{M_\alpha \cap B_{2\delta}} |V \lrcorner \text{Im } \Omega|_{M_\alpha}|_0 \cdot C |d\varphi|_0 \right| + C\delta^{(n-2)/2} \\ &\leq C\delta^{n-1} + C\delta^{(n-2)/2}. \end{aligned}$$

We now consider the second term. Observe that since $\text{Im } \dot{\Omega}$ is a form on X defined independently of α , $|\text{Im } \dot{\Omega}|_{M_\alpha}|_0$ is bounded independently of α . And

obviously $|\operatorname{Im} \dot{\Omega}|_{M_1 \amalg M_2}|_{0, M_1 \amalg M_2}$ is bounded independently of α . Therefore

$$\begin{aligned}
\int_{M_\alpha} (\operatorname{Im} \dot{\Omega}|_{M_\alpha}) S &= \int_{M_\alpha} (\operatorname{Im} \dot{\Omega}|_{M_\alpha}) \varphi \bar{S} + O(\delta^{(n-2)/2}) \text{ by Lemma 11} \\
&= \int_{M_1 \amalg M_2} (\operatorname{Im} \dot{\Omega}|_{M_1 \amalg M_2}) \varphi \bar{S} + O(\delta^{(n-2)/2}) \\
&= \int_{M_1 \amalg M_2} (\operatorname{Im} \dot{\Omega}|_{M_1 \amalg M_2}) \bar{S} + O(\delta^{n/2}) + O(\delta^{(n-2)/2}) \text{ by Lemma 11} \\
&= \sqrt{\frac{\operatorname{Vol}(M_1) \operatorname{Vol}(M_2)}{\operatorname{Vol}(M_1) + \operatorname{Vol}(M_2)}} \left\langle \frac{[M_1]}{\operatorname{Vol}(M_1)} - \frac{[M_2]}{\operatorname{Vol}(M_2)}, [\operatorname{Im} \dot{\Omega}] \right\rangle \\
&\quad + O(\delta^{(n-2)/2}) \text{ by definition of } \bar{S}.
\end{aligned}$$

Now the result follows from Lemma 16. $\square$

6 The Full Linearized Deformation Operator

We are now ready to prove the full injectivity estimate.

Proposition 19 (Full Injectivity Estimate). *For small enough ν independent of α , $\operatorname{DF}_\alpha(0, 0)$ satisfies the injectivity estimate $C_I(\alpha) = C\epsilon^{-\nu}$ for sufficiently small α . That is,*

$$\|\operatorname{DF}_\alpha(0, 0)(u, a)\|_{\mathcal{B}'_\alpha} \geq \frac{1}{C} \epsilon^\nu \|(u, a)\|_{\mathcal{B}_\alpha}.$$

Proof.

$$\begin{aligned}
|a| &\leq C \left| \int_{M_\alpha} a \psi S \right| \text{ by the Key Lemma} \\
&= C \left| \int_{M_\alpha} (\Delta u + a \psi) S \right| \text{ since } \Delta u \text{ is orthogonal to } S \\
&= C \left| \int_{M_\alpha} \rho^2 (\Delta u + a \psi) \rho^{-2} S \right| \\
&\leq C \|\Delta u + a \psi\|_{\mathcal{B}'_\alpha} \int_{M_\alpha} |\rho^{-2} S| \\
&\leq C \|\Delta u + a \psi\|_{\mathcal{B}'_\alpha}
\end{aligned}$$

where the last line follows from the bounds on $|S|_0$ and $\|\rho^{-1}\|_{L^2(M_\alpha)}$. On the other hand,

$$\begin{aligned}
\|u\|_{\mathcal{B}_{1,\alpha}} &\leq C \epsilon^{-\nu} \|\Delta u\|_{\mathcal{B}'_\alpha} \text{ by the Laplacian Injectivity Estimate} \\
&\leq C \epsilon^{-\nu} (\|\Delta u + a \psi\|_{\mathcal{B}'_\alpha} + \|a \psi\|_{\mathcal{B}'_\alpha}) \\
&\leq C \epsilon^{-\nu} \|\Delta u + a \psi\|_{\mathcal{B}'_\alpha}
\end{aligned}$$

where the last line follows from the previous calculation and the fact that $|\psi|_{C_{\rho}^{0,\beta}(M_\alpha)}$ is bounded independently of α , by the definition of ψ . Finally, we deal with the Pu term.

$$\begin{aligned} \|(u, a)\|_{\mathcal{B}_\alpha} &\leq C\epsilon^{-\nu}\|\Delta u + a\psi\|_{\mathcal{B}'_\alpha} \text{ by combining the previous two calculations} \\ &\leq C\epsilon^{-\nu}(\|DF_\alpha(0,0)(u, a)\|_{\mathcal{B}'_\alpha} + \|Pu\|_{\mathcal{B}'_\alpha}) \\ &\leq C\epsilon^{-\nu}\|DF_\alpha(0,0)(u, a)\|_{\mathcal{B}'_\alpha} + C\epsilon^{-\nu}\alpha^{1-\beta}\|(u, a)\|_{\mathcal{B}_\alpha} \text{ by Lemma 9.} \end{aligned}$$

For sufficiently small ν , $\epsilon^{-\nu}\alpha^{1-\beta} \rightarrow 0$ as $\alpha \rightarrow 0$. So for small enough α , we can absorb the last term into the left-hand side. $\square$

Proposition 20. *For small ν , $DF_\alpha(0,0)$ is surjective for sufficiently small α .*

Proof. Consider the map $A : \mathcal{B}_\alpha \rightarrow \mathcal{B}'_\alpha$ defined by $A : (u, a) \mapsto \Delta u + a\psi$. By the proof of the Full Injectivity Estimate together with the Key Lemma, it is evident that A is an isomorphism with $\|A^{-1}\| \leq C\epsilon^{-\nu}$. By Lemma 9, $\|P\| \leq C\alpha^{1-\beta}$, therefore $\|A^{-1}P\| \leq C\alpha^{1-\beta}\epsilon^{-\nu}$. For small enough ν , $\alpha^{1-\beta}\epsilon^{-\nu} \rightarrow 0$ as $\alpha \rightarrow 0$, therefore $I + A^{-1}P$ is invertible, and it follows that $A + P = DF_\alpha(0,0)$ is surjective. $\square$

7 Solving the Deformation Problem

The following Proposition can be found in [16].

Proposition 21 (Nonlinear Estimate). *For small α , F_α satisfies a nonlinear estimate with $C_N = C\epsilon^{-2}$ and $r_1 = \frac{1}{C}\epsilon^2$. That is, for $(h, t) \in \mathcal{B}_\alpha$ with $\|(h, t)\|_{\mathcal{B}_\alpha} \leq r_1$,*

$$\|DF_\alpha(h, t)(u, a) - DF_\alpha(0, 0)(u, a)\|_{\mathcal{B}'_\alpha} \leq C\epsilon^{-2}\|(h, t)\|_{\mathcal{B}_\alpha} \cdot \|(u, a)\|_{\mathcal{B}_\alpha}$$

for all $(u, a) \in \mathcal{B}_\alpha$.

The bound on r_1 is needed so that we can always assume that $\tau = 1$ in our definition of F_α . Finally, we have the following simple estimate.

Proposition 22 (Estimate of $F_\alpha(0,0)$). *For small enough α ,*

$$\|F_\alpha(0,0)\|_{\mathcal{B}'_\alpha} \leq C\alpha^3.$$

Proof. Note that $F_\alpha(0,0) = \langle \text{Im } \Omega, \text{Vol}_{M_\alpha} \rangle_{M_\alpha} = \sin \theta$. Recall that $\rho(x) \leq C\delta$ for $x \in M_\alpha \cap B_\delta$ and $\delta = \frac{\alpha}{C\delta}$. Then since $\sin \theta$ is supported in $M_\alpha \cap B_\delta$,

$$|\rho^2 \sin \theta|_0 \leq C\delta^2 |\sin \theta|_0 \leq C\alpha^3 \quad (10)$$

where the second inequality follows from the bound on $\sin \theta$ from Lemma 6. Now we will estimate $[\rho^{2+\beta} \sin \theta]_\beta$ by interpolation. As in (10), we see that

$$|\rho^{2+\beta} \sin \theta|_0 \leq C\alpha^{3+\beta}.$$

We also have

$$\begin{aligned}
|\nabla(\rho^{2+\beta} \sin \theta)|_0 &\leq |(2+\beta)\rho^{1+\beta}(\nabla \rho) \sin \theta|_0 + |\rho^{2+\beta} \nabla(\sin \theta)|_0 \\
&\leq C\delta^{1+\beta} |\sin \theta|_0 + C\delta^{2+\beta} |\nabla(\sin \theta)|_0 \\
&\leq C\alpha^{2+\beta}
\end{aligned}$$

where the second line follows from the bound on $\nabla \rho$ and the same reasoning as in (10), and the last line uses the bounds on $\sin \theta$ and $\nabla(\sin \theta)$ from Lemma 6. Combining the two previous inequalities, we see that

$$[\rho^{2+\beta} \sin \theta]_\beta \leq C\alpha^3.$$

□

Finally, let $r = (2C_I C_N)^{-1} = \frac{1}{C} \epsilon^{2+\nu}$, which is less than r_1 for sufficiently small α . For small enough ν , we have $\frac{r}{2C_I} = \frac{1}{C} \epsilon^{2+2\nu} > C\alpha^3 \geq \|F_\alpha(0, 0)\|_{\mathcal{B}_\alpha}$ for sufficiently small α . We can now invoke the Inverse Function Theorem to find a solution $F_\alpha(h, t) = 0$ with $\|(h, t)\|_{\mathcal{B}_\alpha} \leq r$, and by elliptic regularity, h is smooth. Since $\|\nabla h\|_0 \leq C\epsilon^{1+\nu}$, it follows that there exists an embedded special Lagrangian submanifold of $(X, J_t, \omega_t, \Omega_t)$ in a $C\epsilon^{1+\nu}$ -neighborhood of M_α for some $t < r$. Finally, since the construction of M_α and F_α can be made to depend smoothly on α , and there is a unique solution to $F_\alpha(h, t) = 0$ in $B_r(0, 0)$, we can also say that the embedded SLags we constructed, as well as t , depend smoothly on α . This concludes the proof of the Main Theorem and the Torus Case.

References

- [1] Adrian Butscher, *Regularizing a singular special Lagrangian variety*, arXiv:math.DG/0110053, to appear in Comm. Anal. Geom.
- [2] ———, *Deformation theory of minimal Lagrangian submanifolds*, Ph.D. thesis, Stanford University, August 2000.
- [3] Philip Candelas and Xenia C. de la Ossa, *Moduli space of Calabi-Yau manifolds*, Strings '90 (College Station, TX, 1990), World Sci. Publishing, River Edge, NJ, 1991, pp. 401–429.
- [4] Jeff Cheeger and Detlef Gromoll, *The splitting theorem for manifolds of nonnegative Ricci curvature*, J. Differential Geometry **6** (1971/72), 119–128.
- [5] Marcos Dajczer, *Submanifolds and isometric immersions*, Mathematics Lecture Series, vol. 13, Publish or Perish Inc., Houston, TX, 1990, Based on the notes prepared by Mauricio Antonucci, Gilvan Oliveira, Paulo Lima-Filho and Rui Tojeiro.
- [6] F. Reese Harvey, *Spinors and calibrations*, Perspectives in Mathematics, vol. 9, Academic Press Inc., Boston, MA, 1990.

- [7] Mark Haskins, *Special Lagrangian cones*, arXiv:math.DG/0005164.
- [8] Nigel J. Hitchin, *The moduli space of special Lagrangian submanifolds*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **25** (1997), no. 3-4, 503–515 (1998), Dedicated to Ennio De Giorgi.
- [9] Dominic Joyce, *Lectures on Calabi-Yau and special Lagrangian geometry*, Part I of M. Gross, D. Huybrechts and D. Joyce, “Calabi-Yau Manifolds and Related Geometries,” Springer, 2003, is an expanded version of this paper, arXiv:math.DG/0108088.
- [10] ———, *Special Lagrangian submanifolds with isolated conical singularities. I. Regularity*, arXiv:math.DG/0211294.
- [11] ———, *Special Lagrangian submanifolds with isolated conical singularities. II. Moduli spaces*, arXiv:math.DG/0211295.
- [12] ———, *Special Lagrangian submanifolds with isolated conical singularities. III. Desingularization, the unobstructed case*, arXiv:math.DG/0302355.
- [13] ———, *Special Lagrangian submanifolds with isolated conical singularities. IV. Desingularization, obstructions and families*, arXiv:math.DG/0302356.
- [14] ———, *Compact manifolds with special holonomy*, Oxford Mathematical Monographs, Oxford University Press, Oxford, 2000.
- [15] Gary Lawlor, *The angle criterion*, Invent. Math. **95** (1989), no. 2, 437–446.
- [16] Yng-Ing Lee, *Embedded special Lagrangian submanifolds in Calabi-Yau manifolds*, to appear in Comm. Anal. Geom.
- [17] Robert C. McLean, *Deformations of calibrated submanifolds*, Comm. Anal. Geom. **6** (1998), no. 4, 705–747.
- [18] Richard M. Schoen, *Special Lagrangian submanifolds*, from the Clay Mathematics Institute Summer School on the Global Theory of Minimal Surfaces.
- [19] Andrew Strominger, Shing-Tung Yau, and Eric Zaslow, *Mirror symmetry is T-duality*, Nuclear Phys. B **479** (1996), no. 1-2, 243–259.
- [20] Gang Tian, *Smoothness of the universal deformation space of compact Calabi-Yau manifolds and its Petersson-Weil metric*, Mathematical aspects of string theory (San Diego, Calif., 1986), Adv. Ser. Math. Phys., vol. 1, World Sci. Publishing, Singapore, 1987, pp. 629–646.
- [21] Andrey N. Todorov, *The Weil-Petersson geometry of the moduli space of $SU(n \geq 3)$ (Calabi-Yau) manifolds. I*, Comm. Math. Phys. **126** (1989), no. 2, 325–346.